# ERRATUM

# Erratum to: Interpersonal discrimination and health-related quality of life among black and white men and women in the United States

Sherrill L. Sellers · Dasha Cherepanov · Janel Hanmer · Dennis G. Fryback · Mari Palta

Published online: 10 January 2013 © Springer Science+Business Media Dordrecht 2013

# Abstract

*Objective* We assessed associations between discrimination and health-related quality of life among black and white men and women in the United States.

*Methods* We examined data from the National Health Measurement Study, a nationally representative sample of 3,648 adults aged 35–89 in the non-institutionalized US population. These data include self-reported lifetime and everyday discrimination as well as several health utility

Erratum to: Qual Life Res DOI 10.1007/s11136-012-0258-8 There was a spelling error in the second author's family name in the original publication. The name is correct in this erratum. In addition, the entire, corrected content of the original article is republished to address multiple errors in the list of references and citations in the text.

The online version of the original article can be found under doi: 10.1007/s11136-012-0258-8.

S. L. Sellers (⊠) Miami University-Ohio, Oxford, OH 45056, USA e-mail: slsellers@muohio.edu

D. Cherepanov Partnership for Health Analytic Research, LLC, Beverly Hills CA, USA e-mail: dgcherepanov@gmail.com

J. Hanmer Department of Internal Medicine, University of Iowa Hospitals and Clinics, Iowa City, IA 52242, USA e-mail: janel-hanmer@uiowa.edu

D. G. Fryback · M. Palta University of Wisconsin-Madison, Madison, WI 53724, USA e-mail: dfryback@wisc.edu

M. Palta e-mail: mpalta@facstaff.wisc.edu indexes (EQ-5D, HUI3, and SF-6D). Multiple regression was used to compute mean health utility scores adjusted for age, income, education, and chronic diseases for each raceby-gender subgroup.

*Results* Black men and women reported more discrimination than white men and women. Health utility tended to be worse as reported discrimination increased. With a few exceptions, differences between mean health utility scores in the lowest and highest discrimination groups exceeded the 0.03 difference generally considered to be a clinically significant difference.

*Conclusions* Persons who experienced discrimination tended to score lower on health utility measures. The study also revealed a complex relationship between experiences of discrimination and race and gender. Because of these differential social and demographic relationships caution is urged when interpreting self-rated health measures in research, clinical, and policy settings.

**Keywords** Health-related quality of life · Interpersonal discrimination · Race and gender differences

# Introduction

Health-related quality of life (HRQL) measures, which include disease specific measures, generic health status measures (e.g., SF-36v2<sup>TM</sup>), and preference-based utility measures (e.g., EQ-5D), are used to evaluate clinical outcomes, determine the burden of disease in a population and quantify health for economic analysis [1]. Studies suggest disparities in HRQL with low scores for those of lower income and education compared with those of higher socioeconomic status, for blacks compared with whites, and for women compared with men [2–5].

Little is known about the mechanisms that underlie these race and gender differences [6, 7]. One factor found to be associated with disparities in health is experience of discrimination. Conceptualized as a social stress, discrimination has been associated with poorer physical and mental health [8, 9]. Yet the prevalence of discrimination across demographic characteristics in the United States is poorly described [10]. In studies using the everyday and lifetime discrimination scales, the most consistent finding is that whites report experiencing less discrimination than nonwhites [9, 10]. Although less consistently found, it appears that prevalence of discrimination may vary by gender [11–13]. Although black men and women were equally likely to perceive general workplace discrimination or mistreatment when shopping in public, [14] black men appear more likely to perceive themselves to face racial discrimination in hiring and in encounters with the police [15].

Few discrimination studies have sufficient sample sizes to simultaneously consider subgroups by race and gender and none have evaluated HRQL. Race and gender are both associated with a range of constrained opportunities and resources, such as differential social capital and exposure to social risks such as discrimination that influence daily life, perceptions, attitudes, health, and well being [16–18]. A more complete understanding of disparities in HRQL requires attention to the intersection of race and gender and experiences of discrimination. We examine our hypothesis that perceived discrimination is associated with lower HRQL scores among black and white men and women from a US sample of adults.

# Methods

# Data

We used cross-sectional data from the US National Health Measurement Study (NHMS) [19] on respondents selfidentifying as either African American/black or white. The NHMS was a random-digit dialed telephone survey, conducted in 2005–2006, of 3,844 community-living US adults aged 35–89 years. The NHMS collected self-reported data on interpersonal discrimination, socioeconomic status, presence of health conditions, and several widely used generic HRQL instruments. NHMS data are publicly available (http://www.icpsr.umich.edu/cocoon/NACDA/ STUDY/23263.xml).

# Health utility measures

This study examined three commonly used utility measures of HRQL: the EuroQol EQ-5D (EQ-5D) [20, 21], the Health Utilities Index Mark 3 (HUI3) [22], and the SF-6D,

computed from the SF-36v2<sup>TM</sup> [23, 24]. The EQ-5D, HUI3, and SF-6D are all generic measures, not specific to any one organ system or disease, and preference-based, their scoring is based on systematically elicited utility evaluations by people sampled from a community or the general population. All three produce summary utility scores anchoring "dead" at 0 and "full health" at 1.0.

The EQ-5D (http://www.euroqol.org) has five questions, each with three response categories: no problem, some problem, or severe problem. The questions refer to "your health today" and ask about mobility, self-care, usual activities, pain/discomfort, and anxiety/depression. We use the scoring algorithm that was derived in a population sample of 4,000 US adults [21].

The HUI3 (http://www.healthutilities.com/) questions refer to "your level of ability or disability during the past week" and access health status on eight attributes: vision, hearing, speech, ambulation, dexterity, emotion, cognition, and pain with five or six levels per attribute, varying from highly impaired to normal [22]. The scoring algorithm for the HUI3 was derived in a community sample from Hamilton, Ontario, Canada.

The SF-36v2<sup>TM</sup> health status questionnaire (http://www.sf-36.org/), consisting of 36 questions generally referring to health in the past 4 weeks, is usually scored as either eight individual scales or two summary scores. Brazier et al. [23, 24] developed the SF-6D, a preference-based scoring algorithm that yields a single summary score using 11 of the items from the SF-36 that defines health status on six attributes: physical function, role limitation, social function, pain, mental health, and vitality. Scoring of the SF-6D is based on a population sample from the United Kingdom [23, 24].

# Discrimination scales

Two scales of interpersonal discrimination were administered to all respondents: the everyday and lifetime discrimination scales (Table 1) [25]. Scores were computed by summing the numerical equivalents for the categorical responses; everyday discrimination scores range from 0 to 25 and lifetime discrimination scores range from 0 to 4, with higher scores representing more discrimination.

# Covariates

The NHMS also surveyed years of education (<12 years, 12 years, >12 years), household income (<\$25,000, \$25,000-\$50,000, \$50,000-\$75,000, and >\$75,000), and past diagnoses of health conditions. We included the five conditions with highest prevalence and impact on HRQL scores: coronary heart disease, arthritis, chronic respiratory disease (asthma, emphysema, and chronic bronchitis), diabetes, and stroke [26].

Everyday discrimination questions (responses: almost everyday = 5, at least once a week = 4, a few times a month = 3, a few times a year = 2, less than once a year = 1, never = 0)

In your day-to-day life, how often have any of the following things happened to you?

- 1. You are treated with less courtesy or respect than other people
- 2. You received poorer service than other people at restaurants or stores
- 3. People act as if they think you are not smart
- 4. People act as if they are afraid of you
- 5. You are threatened or harassed

Lifetime discrimination questions (responses: yes = 1, no = 0)

1. At any time in your life have you ever been unfairly fired or denied promotion?

2. For unfair reasons, have you ever not been hired for a job?

3. Have you ever been unfairly stopped, searched, questioned, physically threatened, or abused by the police?

4. Have you ever been unfairly discouraged by a teacher or advisor from continuing your education?

Table 2 Weighted statistics by race and gender

Item	Black women	White women	Black men	White men	All
Number	705	1,394	381	1,168	3,648
Mean age (year)	50.7	54.2	51.3	55.8	54.5
Lifetime discrimination se	core (%)				
0	41	66	25	55	58
1	31	22	29	28	25
2	18	9	17	12	11
3–4	10	4	29	5	6
Everyday discrimination	score (%)				
0–4	45	66	43	62	62
5–9	36	28	36	30	30
10–14	14	6	14	7	7
15–25	5	1	7	1	1
Household income level (	(%)				
<\$25,000	40	15	29	12	16
\$25,000-\$50,000	25	25	26	25	25
\$50,000-\$75,000	13	21	20	24	22
>\$75,000	22	39	26	39	37
Education level (%)					
<12 year	18	8	11	6	8
=12 year	25	29	30	29	29
>12 year	57	63	60	65	63
Major chronic conditions	(not mutu	ally exclu	isive) (%	<b>b</b> )	
Diabetes	20	9	23	13	12
Coronary heart disease	8	6	11	13	9
Arthritis	33	34	26	30	32
Chronic respiratory disease	20	12	12	13	13
Stroke	6	4	3	3	4

Percentages may not add to 100 % due to rounding

#### Analyses

Means and percentages of discrimination and health utility scores were computed for four race and gender groups. Each race-by-gender group was further stratified by discrimination scores and mean health utility scores were estimated within each stratum using weighted least squares multivariable regression to adjust for age, income, education, and conditions. All adjustment variables were centered to the NHMS weighted means for blacks and whites combined. All analyses were performed in SAS, incorporating survey weights to account for the NHMS sampling design (Copyright 2002–2003 SAS Institute Inc., Cary, NC, USA).

# Results

There were 3,648 respondents: 2,562 white and 1,086 black (Table 2). The sample was predominantly women (57.5 %) with a mean age of 54.5 years. Black men reported the highest lifetime discrimination scores, followed by black women; white women tended to report the least lifetime discrimination. The distribution of everyday discrimination scores was very similar for black men and women and higher than whites' scores. Blacks also had lower income levels than whites. Black men and women reported significantly higher percentages of diabetes than their white counterparts. Black women reported significantly more chronic respiratory disease than other groups.

Health utility scores were worse at higher levels of everyday discrimination in all race-by-gender strata for all HRQL measures except for EQ-5D among black men and

Everyday discrimination score	Blacks and whites		Black men		Black	women	White men		White women		
	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)	
Health utility measured by EQ-	-5D										
0–4	0.88	(0.87, 0.89)	0.89	(0.86, 0.92)	0.85	(0.80, 0.90)	0.88	(0.87, 0.90)	0.88	(0.87, 0.89)	
5–9	0.87	(0.86, 0.88)	0.88	(0.85, 0.91)	0.85	(0.82, 0.88)	0.88	(0.87, 0.90)	0.87	(0.85, 0.88)	
10–25	0.83	(0.80, 0.86)	0.89	(0.83, 0.94)	0.74	(0.69, 0.79)	0.83	(0.79, 0.87)	0.84	(0.80, 0.89)	
Test for trend	<i>p</i> < 0.001		p = 0.23		p = 0.01		p = 0.09		p < 0.001		
Health utility measured by HU	13										
0–4	0.83	(0.81, 0.84)	0.88	(0.85, 0.92)	0.76	(0.68, 0.84)	0.83	(0.80, 0.85)	0.83	(0.81, 0.84)	
5–9	0.79	(0.77, 0.81)	0.82	(0.77, 0.87)	0.78	(0.72, 0.84)	0.81	(0.78, 0.84)	0.78	(0.75, 0.81)	
10–25	0.72	(0.67, 0.77)	0.84	(0.73, 0.96)	0.58	(0.49, 0.66)	0.72	(0.65, 0.79)	0.71	(0.62, 0.80)	
Test for trend	<i>p</i> < 0.0	001	p = 0.25		p = 0.02		p = 0.02		p < 0.001		
Health utility measured by SF-	6D										
0–4	0.80	(0.80, 0.81)	0.83	(0.81, 0.86)	0.79	(0.74, 0.83)	0.81	(0.80, 0.82)	0.80	(0.78, 0.81)	
5–9	0.78	(0.76, 0.79)	0.83	(0.80, 0.85)	0.78	(0.75, 0.81)	0.79	(0.77, 0.80)	0.76	(0.75, 0.78)	
10–25	0.74	(0.72, 0.77)	0.79	(0.73, 0.84)	0.68	(0.63, 0.72)	0.74	(0.71, 0.77)	0.75	(0.71, 0.79)	
Test for trend	p < 0.0	p < 0.001		p = 0.17		p = 0.002		<i>p</i> < 0.001		p < 0.001	

Table 3 Mean health utility score by race and gender on three health utility indexes as a function of everyday discrimination score

Weighted least squares (WLS) multivariable regression was used to compute mean health utility scores adjusted for age, income, education, and chronic diseases within race-by-gender groups stratified by discrimination scale scores. Tests for trend were computed using aforementioned WLS regression models refitted within the four gender-by-race groups (not stratified by discrimination scores); *CI* confidence interval

Table 4	Mean health	utility	score by	race and	gender	on three	health	utility	indexes	as a	a function	of lifetime	discrimination score
---------	-------------	---------	----------	----------	--------	----------	--------	---------	---------	------	------------	-------------	----------------------

Lifetime discrimination score	Blacks and whites		Black men		Black	women	White men		White women	
	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)	Mean	(95 % CI)
Health utility measured by EQ	-5D									
0	0.89	(0.88, 0.89)	0.92	(0.88, 0.96)	0.86	(0.83, 0.89)	0.90	(0.89, 0.91)	0.88	(0.87, 0.89)
1	0.87	(0.86, 0.88)	0.89	(0.86, 0.93)	0.82	(0.73, 0.91)	0.88	(0.85, 0.90)	0.86	(0.84, 0.88)
2	0.84	(0.82, 0.86)	0.90	(0.87, 0.93)	0.82	(0.76, 0.88)	0.84	(0.81, 0.87)	0.85	(0.81, 0.89)
3–4	0.84	(0.80, 0.88)	0.82	(0.78, 0.86)	0.81	(0.75, 0.87)	0.85	(0.77, 0.93)	0.87	(0.79, 0.94)
Test for trend	p < 0.001		p = 0.08		p = 0.01		<i>p</i> < 0.001		p = 0.001	
Health utility measured by HU	JI3									
0	0.83	(0.82, 0.85)	0.91	(0.87, 0.94)	0.80	(0.76, 0.84)	0.85	(0.83, 0.87)	0.83	(0.81, 0.85)
1	0.79	(0.77, 0.81)	0.86	(0.81, 0.91)	0.70	(0.57, 0.84)	0.80	(0.77, 0.83)	0.78	(0.74, 0.81)
2	0.75	(0.72, 0.79)	0.80	(0.72, 0.89)	0.77	(0.69, 0.85)	0.77	(0.73, 0.82)	0.74	(0.68, 0.81)
3–4	0.74	(0.66, 0.82)	0.76	(0.69, 0.83)	0.70	(0.59, 0.82)	0.75	(0.64, 0.86)	0.76	(0.62, 0.90)
Test for trend	p < 0.0	001	p = 0.01		p = 0.02		<i>p</i> < 0.001		p < 0.001	
Health utility measured by SF-	-6D									
0	0.80	(0.80, 0.81)	0.87	(0.85, 0.89)	0.80	(0.77, 0.83)	0.82	(0.80, 0.83)	0.79	(0.78, 0.80)
1	0.78	(0.77, 0.80)	0.85	(0.82, 0.87)	0.75	(0.69, 0.82)	0.79	(0.77, 0.81)	0.77	(0.75, 0.79)
2	0.76	(0.75, 0.78)	0.83	(0.80, 0.85)	0.76	(0.73, 0.80)	0.77	(0.74, 0.79)	0.76	(0.74, 0.79)
3–4	0.75	(0.71, 0.78)	0.76	(0.73, 0.80)	0.67	(0.61, 0.74)	0.75	(0.69, 0.80)	0.78	(0.71, 0.86)
Test for trend	<i>p</i> < 0.001		<i>p</i> < 0.001		<i>p</i> < 0.001		<i>p</i> < 0.001		p = 0.002	

Weighted least squares (WLS) multivariable regression was used to compute mean health utility scores adjusted for age, income, education, and chronic diseases within race-by-gender groups stratified by discrimination scale scores. Tests for trend were computed using aforementioned WLS regression models refitted within the four gender-by-race groups (not stratified by discrimination scores); *CI* confidence interval

white men (Table 3). For SF-6D and HUI3, differences between mean utilities in the lowest and highest discrimination groups exceeded 0.03, a difference considered clinically significant [27, 28].

Higher lifetime discrimination scores were associated with worse health utility scores (Table 4). This difference reached statistical significance (p < 0.05) for SF-6D and HUI3 in all groups, but not for EQ-5D in black men. All differences were deemed clinically significant, except for EQ-5D and SF-6D in white women.

# Discussion

We explored differences in HRQL by discrimination, within race and gender groups. We found higher discrimination was associated with lower health utility scores for blacks and whites, men and women. With a few exceptions, mean health utility score differences between lowest and highest everyday or lifetime discrimination levels in any race-gender strata exceeded 0.03, a clinically significant difference for utility measures. This suggests that the stress of discrimination is pervasive by race and gender [29].

Our study provides nationally representative estimates of both everyday and lifetime discrimination for communitydwelling adults aged 35–89 years. Consistent with other studies, blacks report more discrimination than whites [10]. Black men and women report similar levels of everyday discrimination, yet black men report more lifetime discrimination than black women. White women report the least lifetime discrimination. The high prevalence of discrimination among black men supports a recent theory on racial hierarchy which suggests that subordinate males are primary targets of discrimination and that discrimination is practiced to reduce competition for power [30].

We were unable to assess causation. Poor health may heighten perceptions of unfair treatment [31], although evidence from longitudinal studies indicates that discrimination precedes poor health [32, 33]. Additionally, it is possible that the self-reported survey items in this study were perceived and used differently by race and gender [34–36]. The small sample size for black men is another limitation. Longitudinal studies with diverse population groups are needed to assess causation and generalizability of the study findings.

Race and gender are important and intertwined social constructs that contribute to differences in health, possibly moderated by discrimination [37]. Our study emphasizes the necessity for further studies and development of theory to explain the simultaneous impact of race, gender, and discrimination on health. We urge that extrapolation of health utility differences across these groups in research, clinical, and policy settings should be done with caution.

#### References

- 1. McDowell, I. (2006). *Measuring health: A guide to rating scales and questionnaires*. New York: Oxford University Press.
- Fu, A. Z., & Kattan, M. W. (2006). Racial and ethnic differences in preference-based health status measure. *Current Medical Research and Opinion*, 22(12), 2439–2448.
- Franks, P., Gold, M. R., & Fiscella, K. (2003). Sociodemographics, self-rated health, and mortality in the US. *Social Science and Medicine*, 56(12), 2505–2514.
- 4. Cherepanov, D., Palta, M., Fryback, D. G., & Robert, S. A. (2010). Gender differences in health-related quality-of-life are partly explained by sociodemographic and socioeconomic variation between adult men and women in the US: Evidence from four US nationally representative data sets. *Quality of life research: An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation, 19*(8), 1115–1124.
- Robert, S. A., Cherepanov, D., Palta, M., Cross Dunham, N., Feeny, D., & Fryback, D. G. (2009). Socioeconomic status and age variations in health-related quality of life: Results from the national health measurement study. *The Journals of Gerontology Series B, Psychological Sciences and Social Science, 64*(3), 378–389.
- Sellers, S. L., & Newmann, J. P. (2007). Role of material resources in race and gender differences in mental health. New York, NY: American Sociological Association.
- Pereira, C. C. A., Palta, M., Mullahy, J., & Fryback, D. G. (2011). Race and preference-based health-related quality of life measures in the United States. *Quality of Life Research*, 20(6), 969–978.
- Williams, D. R., & Mohammed, S. (2009). Discrimination and racial disparities in health: Evidence and needed research. J Behavior Medicine, 32, 20–47.
- Sternthal, M. J., Slopen, N., & Williams, D. R. (2011). Racial disparities in health: How much does stress really matter? *Du Bois Review*, 8(1), 95–113.
- Paradies, Y. (2006). A systematic review of empirical research on self-reported racism and health. *International Journal of Epidemiology*, 35, 888–901.
- Krieger, N., & Sidney, S. (1996). Racial discrimination and blood pressure: The CARDIA Study of young black and white adults. *American Journal of Public Health*, 86, 1370–1378.
- Kessler, R. C., Mickelson, K. D., & Williams, D. R. (1999). The prevalence, distribution, and mental health correlates of perceived discrimination in the United States. *Journal of Health Society and Behaviour*, 40, 208–230.
- Guyll, M., Matthews, K. A., & Bromberg, J. T. (2001). Discrimination and unfair treatment: Relationship to cardiovascular reactivity among African American and European American woman. *Health Psychology*, 20, 315–325.
- Carter, R. T. (2007). Racism and psychological and emotional injury: Recognizing and assessing race-based traumatic stress. *The Counseling Psychiatrist*, 35(1), 13–105.
- Barnes, L. L., Mendes De Leon, C. F., Wilson, R. S., Bienias, J. L., Bennett, D. A., & Evans, D. A. (2004). Racial differences in perceived discrimination in a community population of older blacks and whites. *Journal of Aging Health*, *16*, 315–337.
- Williams, D. R., Mohammed, J., Leavell, J., & Collins, C. (2010). Race, socioeconomic status, and health: Complexities, ongoing challenges, and research opportunities. *Annals of the New York Academy of Sciences*, 1186, 69–101.
- 17. Bird, D. C., & Rieker, P. P. (2008). Gender and health: The effects of constrained choices and social policies. New York: Cambridge University Press.
- Barnes, L. L., Mendes de Leon, C. F., Lewis, T. T., et al. (2008). Perceived discrimination and mortality in a population-based study of older adults. *American Journal of Public Health*, 98, 1241–1247.

- Fryback, D. G., Dunham, N. C., Palta, M., Hanmer, J., Buechner, J., Cherepanov, D., et al. (2007). US norms for six generic healthrelated quality-of-life indexes from the national health measurement study. *Medical Care*, 45(12), 1162–1170.
- Brooks, R., Rabin, R., & de Charro, F. (2003). The measurement and valuation of health status using EQ-5D: A European perspective. Dordrecht: Kluwer Academics Publishers.
- Shaw, J. W., Johnson, J. A., & Coons, S. J. (2005). US valuation of the EQ-5D health states: Development and testing of the D1 valuation model. *Medical Care*, 43(3), 203–220.
- Feeny, D., Furlong, W., Torrance, G. W., Goldsmith, C. H., Zhu, Z., DePauw, S., et al. (2002). Multiattribute and single-attribute utility functions for the health utilities index mark 3 system. *Medical Care*, 40(2), 113–128.
- Brazier, J., Roberts, J., & Deverill, M. (2002). The estimation of a preference-based measure of health from the SF-36. *Journal of Health Economics*, 21(2), 271–292.
- Brazier, J. E., & Roberts, J. (2004). The estimation of a preference-based measure of health from the SF-12. *Medical Care*, 42(9), 851–859.
- Williams, D., Yu, Y., Jackson, J., & Anderson, N. (1997). Racial differences in physical and mental health: Socioeconomic status, stress, and discrimination. *Journal of Health Psychology*, 2(3), 335–351.
- Lubetkin, E. I., Jia, H., Franks, P., & Gold, M. R. (2005). Relationship among sociodemographic factors, clinical conditions, and health-related quality of life: Examining the EQ-5D in the US general population. *Quality of Life Research*, 14(10), 2187–2196.
- 27. Walters, S., & Brazier, J. (2005). Comparison of the minimally important difference for two health state utility measures: EQ-5D and SF-6D. *Quality of Life Research*, *14*(6), 1523–1532.
- Guyatt, G. H., Walter, S., & Norman, G. (1987). Measuring change over time: Assessing the usefulness of evaluative instruments. *Journal of Chronic Disease*, 40, 171e8.
- 29. De Vogli, R., Ferrie, J. E., Chandola, T., Kivimaki, M., & Marmot, M. G. (2007). Unfairness and health: Evidence from the

Whitehall II study. *Journal of Epidemiology and Community Health*, 61, 513–518.

- 30. Sidanius, J., & Veniegas, R. C. (2000). Gender and race discrimination: The interactive nature of disadvantage. In S. Oskamp (Ed.), *Reducing prejudice and discrimination: The claremont* symposium on applied social psychology (pp. 47–69). Mahwah, NJ: Lawrence Erlbaum Associates.
- Schulz, A., Gravlee, C., Williams, D., Israel, B., Mentz, G., & Rowe, Z. (2006). Discrimination, symptoms of depression, and self-rated health among African America women in Detroit. Results from a longitudinal analysis. *American Journal of Public Health*, 96(7), 1265–1270.
- 32. Sechrist, G. B., Swim, J. K., & Stangor, C. (2004). When do the stigmatized make attributions to discrimination occurring to the self and others? The roles of self-presentation and need for control. *Journal of Personality and Social Psychology*, 87, 111–122.
- Krieger, N. (1999). Embodying inequality: A review of concepts, measures, and methods for studying health consequences of discrimination. *International Journal of Health Services*, 29, 295–352.
- Fleishman, J., & Lawrence, W. (2003). Demographic variation in SF-12 scores: True differences or differential item functioning? *Medical Care*, 41(7), 75–86.
- Fryback, D. G., Palta, M., Cherepanov, D., Bolt, D., & Kim, J.-S. (2010). Comparison of five health-related quality of life indexes using item response theory analysis. *Medical Decision Making*, 30(1), 5–15.
- Lewis, T. T., Yang, M. F., Jacobs, E. A., & Fittchett, G. (2012). Racial/ethnic differences in responses to the everyday discrimination scale: A differential item functioning analysis. *American Journal of Epidemiology*, 175(5), 391–401.
- Hankivsky, O. (2012). Women's health, men's health, and gender and health: Implications of intersectionality. *Social Science & Medicine*, (74)1712–1720.